



## Dynamics of Prolyl hydroxylases levels during disease progression in experimental colitis

Bakshi, H. A., Mishra, V., Satija, S., Mehta, M., Hakkim, F. L., Kesharwani, P., Dua, K., Chellappan, D. K., Charbe, N. B., Shrivastava, G., Rajeshkumar, S., Aljabali, A. A., Al-trad, B., Pabreja, K., & Tambuwala, M. M. (2019). Dynamics of Prolyl hydroxylases levels during disease progression in experimental colitis. *Inflammation*, 42(6), 2032-2036. <https://doi.org/10.1007/s10753-019-01065-3>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Inflammation

**Publication Status:**  
Published (in print/issue): 01/12/2019

**DOI:**  
[10.1007/s10753-019-01065-3](https://doi.org/10.1007/s10753-019-01065-3)

**Document Version**  
Author Accepted version

### General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).

## Dynamics of prolyl hydroxylases levels during disease progression in experimental colitis

Hamid A. Bakshi<sup>1</sup>, Vijay Mishra<sup>2</sup>, Saurabh Satija<sup>2</sup>, Meenu Metha<sup>2</sup>, Faruk L. Hakkim<sup>3</sup>, Prashant Kesharwani<sup>4</sup>, Kamal Dua<sup>5</sup>, Dinesh K. Chellappan<sup>6</sup>, Nitin B. Charbe<sup>7,8</sup>, Garima Shrivastava<sup>9</sup>, Rajeshkumar S<sup>10</sup>, Alaa A. Aljabali<sup>11</sup>, Bahaa Al –Trad<sup>12</sup>, Kavita Pabreja<sup>13</sup> and Murtaza M. Tambuwala<sup>1\*</sup>

<sup>1</sup>SAAD Centre for Pharmacy and Diabetes, School of Pharmacy and Pharmaceutical Science, Ulster University, Coleraine, County Londonderry, BT52 1SA, Northern Ireland. United Kingdom.

<sup>2</sup>School of Pharmaceutical Sciences, Lovely Professional University, Phagwara (Punjab), India.

<sup>3</sup>Department of Mathematics and Sciences, College of Arts and Applied Sciences, Dhofar University, Salalah, Oman.

<sup>4</sup>School of Pharmaceutical Education and Research, Jamia Hamdard (Hamdard University), New Delhi 110062, India.

<sup>5</sup>Discipline of Pharmacy, Graduate School of Health, University of Technology Sydney, Sydney, NSW 2007, Australia.

<sup>6</sup>Department of Life Sciences, School of Pharmacy, International Medical University, Bukit Jalil 57000, Kuala Lumpur, Malaysia.

<sup>7</sup>Departamento de Química Orgánica, Facultad de Química y de Farmacia, Pontificia Universidad Católica de Chile, Av. Vicuña McKenna 4860, Macul, Santiago 7820436, Chile.

<sup>8</sup>Sri Adichunchunagiri University, Sri Adichunchunagiri College of Pharmacy, BG Nagar, Karnataka 571418, India.

<sup>9</sup>Indian Institute of Technology, Delhi, India.

<sup>10</sup>Department of Pharmacology, Saveetha Dental College and Hospitals, SIMATS, Chennai 600077, TN, India.

<sup>11</sup>Faculty of Pharmacy, Department of Pharmaceutical Sciences, Yarmouk University, Irbid, Jordan.

<sup>12</sup>Department of Biological Sciences, Yarmouk University, 21163 Irbid, Jordan.

<sup>13</sup>School of Medicine and Public Health, University of Newcastle, Newcastle, Australia.

### \*Corresponding author

Dr. Murtaza M. Tambuwala  
SAAD Centre for Pharmacy and Diabetes,  
School of Pharmacy and Pharmaceutical Sciences,  
Ulster University,  
Coleraine, BT52 1SA,  
Northern Ireland.  
United Kingdom.  
E-Mail: m.tambuwala@ulster.ac.uk  
Tel: +44 28-701-24016  
Fax: +44 28-701-23518

**Abstract-** Hypoxia inducible factor (HIF)-prolyl hydroxylase (PHD) inhibitors are shown to be protective in several models of inflammatory bowel disease (IBD). However, these non-selective inhibitors are known to inhibit all the three isoforms of PHD *i.e.* PHD-1, PHD-2 and PHD-3. In the present report, we investigated the associated changes in levels of PHDs during the development and recovery of chemically induced colitis in mice. The results indicated that in the experimental model of murine colitis, levels of both, PHD-1 and PHD-2 were found to be increased with progression of the disease; however, the level of PHD-3 remained the same in group of healthy controls and mice with colitis. Thus, the findings advocated that inhibitors, which inhibited all three isoforms of PHD could not be ideal therapeutics for IBD since PHD-3 is required for normal gut function. Hence this necessitates the development of new compounds capable of selectively inhibiting PHD-1 and PHD-2 for effective treatment of IBD.

**Keywords:** Prolyl hydroxylases, Inflammatory bowel disease, Colitis, Disease activity index

## INTRODUCTION

Hypoxia inducible factor (HIF)-prolyl hydroxylases (PHDs) are member of dioxygenase enzymes family. These enzymes are known to play an important function in intracellular oxygen sensing as well as signalling responses in low oxygen levels (hypoxia). This occurs mainly by the regulation of HIF stability [1-4]. These PHDs consist of three isoforms (PHD-1, PHD-2 and PHD-3), which share mostly similar biochemical characteristics, however all three isoforms of PHDs are known to have very distinct and tissue-specific expression profiles and, regulation of HIF, which in turn can affect pro-survival signalling pathways, including nuclear factor  $\kappa$ B pathways during inflammation [1, 5-7]. These PHDs are known to mediate their effect on HIF pathway via hydroxylation of proline residues of the HIF $\alpha$  subunit, which is degraded under normoxic condition. Once it is hydroxylated, HIF $\alpha$  acts as a target for ubiquitylation by von Hippel Lindau E3 ubiquitin ligase and results in the degradation and ubiquitination of HIF $\alpha$  [8]. The PHDs perform a pivotal function in adaptive response known to occur during hypoxia via the activation of several pathways of genes expression implicated in support of cell survival, erythropoiesis, angiogenesis and metabolism. These features make PHD an interesting target for therapy [9].

Inflammatory bowel disease (IBD), an idiopathic disease with genetically heterogenous nature, has chronic inflammatory conditions in the gastrointestinal tract (GIT) with severe pathology and limited therapeutic options [10-12]. Till date the underlying causes of IBD are not well defined, however the fundamental defect in this disease involves damage/injury of the intestinal epithelial barrier resulting in the development and progression of the disease. Hence targeting of the HIF-PHD

1 pathway may be a fascinating area for potential future therapeutics for inflammatory  
2 disease.

3  
4 Various reports employing pharmacological inhibition of hydroxylase enzyme  
5 revealed improvisation in variety of inflammatory conditions including IBD [13, 15].  
6  
7 Several researchers reported that pan-hydroxylase inhibitors such as  
8 dimethyloxalylglycine (DMOG) have a protective effect in experimental colitis [13-15].  
9  
10 However, majority of these studies have not identified the actual isoform of PHDs or  
11 the effector pathways involved in this protective effect.  
12  
13  
14  
15  
16  
17

18  
19 Our previous findings suggested that there was increased PHD-1 level in  
20 patients with active ulcerative colitis and genetic loss of PHD-1 was found to be  
21 protective in experimental colitis [16]. However, since there are three isoforms of PHDs  
22 it would be of interest for researchers to know how the levels of PHDs change during  
23 disease development and recovery in IBD. Thus to answer this query, the present  
24 study has been framed to examine the changes in levels of PHDs (PHD-1, -2 and -3)  
25 during the inflammation development and its progression in experimental model of  
26 murine colitis.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

## 38 **MATERIALS AND METHODS**

### 39 **Dextran Sodium Sulphate (DSS) Induced Colitis**

40  
41 Colitis was induced in female C57/Bl6 mice using our previously established  
42 protocol [15, 17]. Briefly, 2.5% w/v dextran sodium sulphate (DSS) was administered  
43 in mice via drinking water to induce colitis. Colon tissues were taken at days 2, 3, 4,  
44 5, 6 and 7 from mice exposed to 2.5% w/v DSS in drinking water. The mice of recovery  
45 group were exposed to 2.5% w/v DSS in drinking water for 5 days and allowed to  
46 recover naturally for next 5 days. On the 10<sup>th</sup> day, colon tissue was collected from this  
47 group.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Assessment of Prolyl Hydroxylases (PHD) Levels

The colon tissue lysates were analysed for the level of protein expression of PHD-1, -2 and -3 levels by Western blotting technique as reported previously [16].

## Disease Activity Index (DAI)

The weight of each animal was recorded on a daily basis. The stool consistency and appearance of faecal blood was monitored. These parameters were converted in to Disease Activity Index (DAI) as illustrated previously [18].

## Colon Length

Each excised colon length was recorded after removal of the faecal matter using a cotton swap dipped in phosphate buffer saline (PBS) solution following previously reported method [15, 18].

## Statistical Analysis

Analysis of variance (ANOVA) or Student *t* test was employed for statistical comparisons and data was represented as mean±standard deviation (mean±SD).

## RESULTS AND DISCUSSION

The DAI scores indicated that administration of DSS in drinking water caused weight loss, diarrhoea and appearance of blood in faeces, since the DAI steadily raised from day 2 onwards indicating the onset of inflammation from day 2 (Fig. 1a). We have also noted that colon length of mice progressively decreased as the inflammation progressed (Fig. 1b). This could be due to muscle wasting of the colon tissue during inflammation. Furthermore, both the DAI score and colon length indicated that withdrawal of DSS from drinking water resulted in tissue recovery and alleviation of the classical symptoms of colitis (Fig. 1).

Our key results indicated that PHD-1 level was found to be lower than that of the healthy control at day 2 and it gradually increased by day 5 (Fig. 2). This event

1 suggested that PHD-1 expression levels initially decreased when inflammation sets in  
2 and then progressively increased as the inflammation was established due to  
3 administration of DSS in drinking water. The recovery group has shown the highest  
4 level of PHD-1 expression in colon tissue (Fig. 2a), suggesting that once the PHD-1  
5 level is elevated due to inflammation it does not go down even after the withdrawal of  
6 DSS. The PHD-2 levels followed a similar decline trend at day 2 followed by an  
7 increase at day 5. In case of PHD-2, the recovery group also showed the highest levels  
8 in the colonic tissue (Fig. 2b). Contrary to this, the levels of PHD-3 were observed to  
9 be constant in healthy control, recovery group and during the course of disease  
10 progression (Fig. 2c). It suggested that PHD-3 has not involved in development and  
11 progression of inflammation during experimental colitis.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25  
26 The upregulated PHD-1 in colonic biopsies of IBD patients illustrated the  
27 correlation between PHD1 expression and disease severity in patients [16, 19].  
28 Among PHD-1, -2 and -3, only PHD-1 deficient mice were found to be selectively  
29 protected against DSS-induced colitis development advocating an affirmative function  
30 of PHD-1 in the management of intestinal epithelial cell apoptosis and preservation of  
31 epithelial barrier during intestinal inflammation [16]. The hypothesized link between  
32 improved barrier function and consequent protection against the colitis development  
33 was also supported by studies performed in other models indicating an antiapoptotic  
34 effect of pharmacologically inhibited hydroxylase enzyme [20-22]  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

## 58 CONCLUSION

59  
60  
61  
62  
63  
64  
65

These results suggest that PHD-3 isoform is expressed in healthy as well as diseased colon tissue, indicating that PHD-3 is required for the normal function of the gut. However, PHD-1 and -2 expression levels fall when inflammation is initiated and gradually increases during progression of inflammation and is observed to be the highest in chronic stages. The recovery group indicates that once the agent causing inflammation of the colon *i.e.* DSS is removed from the drinking water, these mice show signs of recovery (Fig. 1), however the increased levels of PHD-1 and -2 are not reduced to normal level, which suggest that PHD-1 and -2 are involved in chronic inflammatory stages of the disease. These findings advocate that inhibition of all PHDs using a non-selective prolyl hydroxylases inhibitor may not be the best strategy towards the development of new therapeutics for IBD. Hence strategies to selectively target PHD-1 and -2 are warranted leading to an improved therapeutic advancement in the field of IBD treatment.

**COMPLIANCE WITH ETHICAL STANDARDS:** All animal experiments were performed under UK Home Office personal and project licence

**Conflict of interest-** Authors declare no conflict of interest.

**Figure legends**



**Fig. 1.** Changes in disease activity score (DAI) and colon length during active colitis.

In **a**, the composite score of weight loss, stool consistency and blood in faeces during disease progression was represented, and in **b**, the changes of colon length during active colitis were given. \*Each control and experimental group contained a minimum of n=6 mice.

**Fig. 2.** Colon homogenates assessed for (a) PHD-1, (b) PHD-2 and (c) PHD-3 protein levels. Each control and experimental group contained a minimum of n=6 mice. PHD-1 level was found to be increased with disease progression in murine model of colitis.

## REFERENCES

1. Kaelin W, Ratcliffe P. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Molecular cell* 2008;30:393-402.
2. Schipani E, Maes C, Carmeliet G, et al. Regulation of Osteogenesis Angiogenesis Coupling by HIFs and VEGF. *Journal of Bone and Mineral Research* 2009;24:1347-1353.
3. Semenza G. Involvement of oxygen-sensing pathways in physiologic and pathologic erythropoiesis. *Blood* 2009;114:2015.
4. Semenza G. Hypoxia inducible factor 1 and cancer pathogenesis. *IUBMB life* 2008;60:591-597.
5. Cummins EP, Berra E, Comerford KM, et al. Prolyl hydroxylase-1 negatively regulates I $\kappa$ B kinase-beta, giving insight into hypoxia-induced NF $\kappa$ B activity. *Proc Natl Acad Sci U S A* 2006;103:18154-9.
6. Chan D, Kawahara T, Sutphin P, et al. Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. *Cancer Cell* 2009;15:527-538.
7. Fu J, Taubman MB. Prolyl hydroxylase EGLN3 regulates skeletal myoblast differentiation through an NF- $\kappa$ B-dependent pathway. *J Biol Chem* 2010;285:8927-35.
8. Oliver KM, Taylor CT, Cummins EP. Hypoxia. Regulation of NF $\kappa$ B signalling during inflammation: the role of hydroxylases. *Arthritis Res Ther* 2009;11:215.
9. Fraisl P, Aragonés J, Carmeliet P. Inhibition of oxygen sensors as a therapeutic strategy for ischaemic and inflammatory disease. *Nature Reviews Drug Discovery* 2009;8:139-152.
10. Braus N, Elliott D. Advances in the pathogenesis and treatment of IBD. *Clinical Immunology* 2009;132:1-9.

11. Rutgeerts P, Vermeire S, Van Assche G. Biological therapies for inflammatory bowel diseases. *Gastroenterology* 2009;136:1182-1197.
12. Imielinski M, Baldassano R, Griffiths A, et al. Common variants at five new loci associated with early-onset inflammatory bowel disease. *Nature genetics* 2009.
13. Cummins E, Seeballuck F, Keely S, et al. The hydroxylase inhibitor dimethyloxalylglycine is protective in a murine model of colitis. *Gastroenterology* 2008;134:346-348.
14. Robinson A, Keely S, Karhausen J, et al. Mucosal Protection by Hypoxia-Inducible Factor (HIF) Prolyl Hydroxylase Inhibition. *Gastroenterology* 2008;134:145.
15. Tambuwala MM, Manresa MC, Cummins EP, et al. Targeted delivery of the hydroxylase inhibitor DMOG provides enhanced efficacy with reduced systemic exposure in a murine model of colitis. *J Control Release* 2015;217:221-7.
16. Tambuwala MM, Cummins EP, Lenihan CR, et al. Loss of prolyl hydroxylase-1 protects against colitis through reduced epithelial cell apoptosis and increased barrier function. *Gastroenterology* 2010;139:2093-101.
17. Khan MN, Lane ME, McCarron PA, et al. Caffeic acid phenethyl ester is protective in experimental ulcerative colitis via reduction in levels of pro-inflammatory mediators and enhancement of epithelial barrier function. *Inflammopharmacology* 2017.
18. Cooper HS, Murthy SN, Shah RS, et al. Clinicopathologic study of dextran sulfate sodium experimental murine colitis. *Lab Invest* 1993;69:238-49.
19. Van Welden, S. et al. (2013) Differential expression of prolyl hydroxylase 1 in patients with ulcerative colitis versus patients with Crohn's disease/infectious colitis and healthy controls. *J. Inflamm. (Lond)* 10, 36.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
20. Liu XB, Wang JA, Ogle ME, et al. Prolyl hydroxylase inhibitor dimethyloxallylglycine enhances mesenchymal stem cell survival. *J Cell Biochem* 2009;106:903–911.
21. Lomb DJ, Straub JA, Freeman RS. Prolyl hydroxylase inhibitors delay neuronal cell death caused by trophic factor deprivation. *J Neurochem* 2007;103:1897–1906.
22. Nangaku M, Izuhara Y, Takizawa S, et al. A novel class of prolyl hydroxylase inhibitors induces angiogenesis and exerts organ protection against ischemia. *Arterioscler Thromb Vasc Biol* 2007; 27:2548–2554.




Figure 2

